

# Astrometric Measurements of Double Star WDS 06047-4505



K. E. Kidwell, L. D. Parsons, M. F. Kidd

Department of Physics, Tennessee Technological University

INSTITUTE FOR STUDENT ASTRONOMICAL RESEARCH

## INTRODUCTION

The research objective was to take astrometric measurements to determine the position angle ( $\theta$ ) and separation ( $\rho$ ) of the AB component of the quintuple system WDS 06047-4505. The Washington Double Star Catalog (WDS) and the Stelle Doppie website were used to determine candidate star systems. Telescopic capabilities (discussed below) and optimal viewing based on the time of year were factors considered in deciding upon criteria of a separation of 5-15 arc seconds and a magnitude between 6-11 for each star. Physical and uncertain doubles observed no later than 2015 that met those criteria were considered and WDS 06047-4505 HJ 3834AB was selected.

The quintuple star system WDS 06047-4505 was originally discovered by John Herschel in 1837 (Sordiglioni, 2015). According to Stelle Doppie, WDS 06047-4505 HJ 3834AB is an “uncertain” double (Sordiglioni, 2015). It has had 42 observations between 1837 and 2015. It was first measured with a position angle of  $246^\circ$  and a separation of  $1.1''$  in 1837. Its most recent measurement was a position angle of  $215^\circ$  and a separation of  $6.0''$  in 2015. The magnitudes of the primary and secondary stars are 6.02 and 8.98, respectively resulting in a delta magnitude ( $\Delta M$ ) of 2.96.

## METHOD

Looking at the calculated orbit of WDS 06047-4505, the most recent observation deviated from the calculated orbit, making this a system of interest. The observation portal was utilized to request images and then twenty images at just under one second exposure time were taken on October 2nd, 4th and 11th, 2020 with the SBIG STL-6303 camera at the Siding Spring, Australia, LCO telescope site. Historical data was obtained from Dr. Brian Mason at the U.S. Naval Observatory as well as from Stelle Doppie.

## RESULTS

Figure 1 shows an observation of WDS 06047-4505 HJ 3834AB taken from the Siding Spring LCO observatory being measured with AstroImageJ software. The primary star is the object in the center and the secondary star is slightly bottom left of the primary.




Figure 1: An image of WDS 06047-4505 taken with the Las Cumbres Observatory 0.4-meter telescope at Siding Spring Observatory

Table 1: Separation and position angle measurements for WDS 06047-4505 HJ 3834AB.

Obs #	Separation (arcsec)	Position Angle (deg)	Exposure Time (sec)	BJD
1	5.82	213.19	0.79	2459125.13
2	5.76	213.52	0.79	2459125.13
3	5.67	211.67	0.79	2459125.13
4	5.79	213.26	0.68	2459127.12
5	5.76	214.89	0.69	2459127.12
6	5.62	215.61	0.69	2459127.12
7	5.68	214.88	0.69	2459127.12
8	5.70	213.53	0.69	2459127.12
9	5.83	213.38	0.69	2459127.12
10	5.72	213.63	0.69	2459127.12
11	5.76	213.54	0.69	2459127.12
12	5.65	213.03	0.68	2459134.11
13	5.97	213.13	0.68	2459134.10
14	5.85	213.52	0.69	2459134.10
15	5.74	213.14	0.68	2459134.10
16	5.76	213.80	0.69	2459134.10
17	5.94	212.76	0.68	2459134.10
18	5.81	213.20	0.69	2459134.10
19	5.88	213.79	0.69	2459134.10
20	5.76	212.93	0.68	2459134.10
Mean	5.77	213.52		
Stan.				
Dev.	0.09	0.84		
SEM	0.02	0.19		

Figure 2 shows the historical data and current measurement plotted on Cartesian coordinates, using Richard Harshaw's Plot Tool (Harshaw, 2020). The original data point from John Herschel was omitted as it was clearly an outlier (shown by a red circle on the cartesian plot). The graph indicates this is a short arc binary system, with an  $R^2$  value of 0.9051.




Figure 2: Plot of historical data and current observation, marked with a red triangle.

Table 2 includes data obtained from the SIMBAD astronomical database, including parallax and proper motion values for the primary and secondary stars. The parallax values are quite similar, 36.6 mas and 37.2 mas, indicating that the stars are very close to each other, and their proper motions in both right ascension (-80.3 and -83.5) and declination (254 and 236 mas/yr) are also similar, suggesting the pair is moving together.

Table 2: Gaia Data

	Primary	Secondary
Parallax (mas)	36.5892	37.2282
Parallax error (mas)	0.2665	0.0607
pmra (mas/yr)	-80.313	-83.454
pmdec (mas/yr)	254.158	236.476
magnitude	6.02	8.98

The current data point lies slightly off the calculated orbit as shown by the red triangle in Figure 3 below. However, it aligns closely with the several most recent speckle interferometry observations. Because speckle observations are generally highly reliable, it may be that a new orbit needs to be calculated based on the past few years of observations.




Figure 3: Orbital diagram of WDS 06047-4505 with the current observation added and marked with a red triangle. The current observation aligns with recent speckle interferometry measurements.

## DISCUSSION

In an effort to distinguish WDS 06047-4505 as either a binary system or an optical double, the historical data was plotted. The cartesian plot of the historical data for WDS 06047-4505 shows the beginning of an arc, indicating this may be a true binary system. One of the challenges analyzing the WDS 06047-4505 system was the large delta magnitude and their close proximity.

Selecting an appropriate aperture was difficult because the star images were so different in size. The quality and resolution of the separation resulted in the need for additional images with shorter exposure times. Comparing the two stars, the primary has a parallax of 36.6 milliarcseconds (mas) with a mean error of 36.3 mas to 36.9 mas while the secondary has 37.2 mas with an error of 37.2 mas to 37.3 mas. Using the weighted parallax method (Harshaw, 2018), it is calculated that the two stars have a weighted minimum separation of 157 AU. The estimation of the physical distance shows a high possibility of the stars being gravitationally bound because “very few known binaries have separations that exceed 3,000 AU, and most are closer than 1,000 AU” (Harshaw, 2018). The calculated value of 157 AU is well within this separation range. Furthermore, the proper motion data from Gaia indicates that the stars are moving at similar rates, in similar directions. Referring to Table 2, the right ascension proper motion (pmra) of the primary star is -80.3 mas/year (yr) and the secondary star is -83.5 mas/yr. They are very large and also very similar in value. The declination proper motion (pmdec) of the primary star is 254 mas/yr while the secondary star has a pmdec of 236 mas/yr. Similarly, the values do not overlap, but they are very close. According to Harshaw (2018), “Two stars that are in orbit around one another should have identical, or very nearly identical, proper motions”. Taken together, the changes in position angle and separation, along with parallax and proper motion data all suggest that this is a gravitationally bound system.

## CONCLUSION

The latest observation of the star WDS 06047-4505 HJ 3834AB yielded a position angle of  $5.77''$  and a separation of  $213.52^\circ$ . A plot of the current observation alongside the historical data on a cartesian graph indicates a short-arc binary with an  $R^2$  value of 0.9051. Three distinct pieces of evidence contribute to the conclusion: the historical evidence, the current measurements, and the data available from Gaia regarding parallax and proper motion. Considering these three pieces of evidence, this star system appears to be a binary star system. This is a promising candidate worthy of further study to confirm that it is in fact a binary system.

## ACKNOWLEDGEMENTS

We would like to thank Rachel Freed and Richard Harshaw for their help with the research. We would also like to thank Dr. Brian Mason for providing historical data. This work has made use of data from the European Space Agency (ESA) mission Gaia (<https://www.cosmos.esa.int/gaia>), processed by the Gaia Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement. It also has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

## REFERENCES

- Harshaw, R. (2018). Gaia DR2 and the Washington Double Star Catalog: A Tale of Two Databases. *JDSO* Vol 14 No 4.
- Harshaw, R. (2020). Using Plot Tool 3.19 to Generate Graphical Representations of the Historical Measurement Data. *JDSO* Vol 16 No 4.
- Sordiglioni, G. (2015). 06047-4505 HJ 3834AB. Retrieved October 29, 2020, from <https://www.stelldoppie.it/index2.php?iddoppia=23095>

## ADDITIONAL CONTRIBUTORS

Kyra Bettwy<sup>2</sup>  
Rolf Hirschfeld<sup>4</sup>  
David Pham<sup>1</sup>  
Corin Slown<sup>3</sup>

- Crean Lutheran High School, CA
- Marymount School of New York, NY
- California State University Monterey Bay, CA
- Kopernikus Foundation, Chile